

CHERN-SIMONS THEORY ON THE TORUS¹

F.Falceto², K.Gawędzki

IHES, Bures-sur-Yvette, 91440, France

Recently a considerable effort has been invested in understanding Chern-Simons theories from the canonical or covariant points of view. The covariant quantization was used to obtain topological invariants of three dimensional manifolds [1] whereas the canonical point of view [1-5] allowed to relate the states of three dimensional Chern-Simons theory to conformal blocks of two dimensional Wess-Zumino-Witten models.

Here we shall discuss the second approach. More precisely we shall study the space of quantum states of the Chern-Simons theory for the group $SU(2)$ on $\mathbb{T}^2 \times \mathbb{R}$ and in the presence of Wilson lines $\{z_n\} \times \mathbb{R}$, $n = 1, \dots, N$, corresponding to representations ρ_{j_n} of spin j_n (acting in spaces V_{j_n}).

The gauge freedom is partially fixed by setting the temporal ($\mathbb{R}$ -direction) component of the connection to zero. The remaining gauge invariance is imposed on the states (very much like the Gauss law in QED) in terms of a quantum flatness condition. The presence of the Wilson lines contributes source terms to the flatness condition.

In the holomorphic quantization *à la Bargmann* we pick up a complex structure on the torus $\mathbb{T}^2$ by fixing the modular parameter $\tau = \tau_1 + i\tau_2 \in \mathbb{H}$. This induces a complex structure in the space of two dimensional (smooth) connections. Integration of the flatness condition leads then to the following picture of the quantum states (see [2], [4], [6],.): they are holomorphic functionals

$$\Psi : \mathcal{A}^{01} \equiv \Omega^{01}(T^2) \otimes \mathfrak{sl}(2, \mathbb{C}) \rightarrow \bigotimes_n V_{j_n}$$

satisfying

$$\Psi({}^h A^{01}) = e^{k S_{WZW}({}^{h^{-1}} A^{01})} \prod_n h(z_n)_{(n)} \Psi(A^{01}), \quad (1)$$

where S_{WZW} is the Wess-Zumino-Witten action coupled to the $(0, 1)$ part of the connection whose gauge transformed version ${}^h A^{01}$ is given by

$${}^h A^{01} = h A^{01} h^{-1} + h \bar{\partial} h^{-1} \quad (2)$$

with h in the gauge group $\mathcal{G}^{\mathbb{C}} \equiv C^\infty(\mathbb{T}^2, \mathrm{SL}(2, \mathbb{C}))$. $h(z)_{(n)} \equiv I \otimes \dots \otimes \rho_{j_n}(h(z)) \otimes \dots \otimes I$.

From (1) it follows that Ψ is determined from its value at a point of each gauge orbit. However the space $\mathcal{A}^{01}/\mathcal{G}^{\mathbb{C}}$ is not a manifold and to study the smoothness of Ψ subject to (1) we need a detailed knowledge of positions and sizes (codimensions) of the $\mathcal{G}^{\mathbb{C}}$ orbits.

First we shall identify $\mathcal{A}^{01}$ with the space of structures of holomorphic $SL(2, \mathbb{C})$ bundles on the trivial bundle $\mathbb{T}^2 \times \mathbb{C}^2$: holomorphic sections of the bundle corresponding to A^{01} are given by maps g with $(\bar{\partial} + A^{01})g = 0$. This space has been studied in [7] and from there we infer that in our case almost every bundle, except for some submanifolds of complex codimension at least 2, is semistable and all these strata are formed by the union of gauge orbits. From the properties of the stratification and the Hartogs theorem, it follows that

¹talk given by the first author at the XIXth International Colloquium on Group Theoretical Methods in Physics, Salamanca (Spain), June 29-July 4, 1992

²Address after October 1st, 1992: Depto. Física Teórica, U. Zaragoza, 50009 Zaragoza, Spain

every holomorphic functional on the space of semistable bundles can be uniquely extended to the whole space.

Every semistable connection can be gauge transformed into one of the following [8][6]

$$\begin{aligned} A_u^{01} &= -u\sigma_3 d\bar{z}/(2\tau_2) & u \in \mathbb{C} \\ \hat{A}_\alpha^{01} &= h_\alpha(-\sigma_+ d\bar{z}/(2\tau_2)) & \alpha = 0, \frac{1}{2}, \frac{\tau}{2}, \frac{\tau+1}{2}, \end{aligned}$$

with $h_\alpha = \exp[(\alpha\bar{z} - \bar{\alpha}z)\sigma_3/(2\tau_2)]$. Note that for $\alpha = 1/2, \tau/2, (\tau+1)/2$, $h_\alpha \notin \mathcal{G}^{\mathbb{C}}$, however its gauge action on $\mathcal{A}^{01}$ is still well defined by (2).

The set $\mathcal{A}_0^{01}$, formed by the union of $\mathcal{G}^{\mathbb{C}}$ orbits of A_u^{01} for $u \in \mathbb{C} \setminus (\mathbb{Z} + \tau\mathbb{Z})/2$, is open dense in $\mathcal{A}^{01}$, while orbits $\mathcal{A}_{(\alpha,0)}^{01} \equiv {}^{\mathcal{G}^{\mathbb{C}}} \hat{A}_\alpha^{01}$ and $\mathcal{A}_{(\alpha,1)}^{01} \equiv {}^{\mathcal{G}^{\mathbb{C}}} A_\alpha^{01}$ are for $\alpha = 0, 1/2, \tau/2, (\tau+1)/2$ complex submanifolds of codimension 1 and 3 respectively (see [6] for details and proofs).

From every holomorphic functional Ψ on $\mathcal{A}^{01}$ we define a holomorphic function $\gamma : \mathbb{C} \rightarrow \otimes_n V_{j_n}$ by

$$\Psi(A_u^{01}) \equiv e^{\pi k u^2 / \tau_2} \prod_n (e^{\sigma_3(\bar{z}_n - z_n)u/(2\tau_2)})_{(n)} \gamma(u). \quad (3)$$

If Ψ satisfies (1) then

$$\gamma(u+1) = \gamma(u), \quad (4,a)$$

$$\gamma(u+\tau) = e^{-2\pi i k(\tau+2u)} \prod_l (e^{iz_n \sigma_3})_{(n)} \gamma(u), \quad (4,b)$$

$$0 = \sum_n (\sigma_3)_{(n)} \gamma(u), \quad (4,c)$$

$$\gamma(-u) = \prod_n (\Omega)_{(n)} \gamma(u), \quad (4,d)$$

with Ω the generator of the Weyl group. These properties come from (1) specialized to the gauge transformations that send connections A_u^{01} to $A_{u'}^{01}$. Conversely, every holomorphic map γ on $\mathbb{C}$ satisfying (4,a-d), defines by (1) and (3) a unique functional Ψ on the open dense stratum $\mathcal{A}_0^{01}$ but this **need not** extend to a holomorphic functional on the whole of $\mathcal{A}^{01}$. We explain now the necessary and sufficient conditions for the existence of such an extension.

Define $g_u = \exp(\frac{1}{2u}\sigma_+)$. Then

$$h_\alpha M_u d\bar{z} \equiv h_\alpha g_u A_u^{01} \xrightarrow{u \rightarrow 0} \hat{A}_\alpha^{01}.$$

From now on we will use a realization of spaces V_{j_n} in terms of polynomials of degree at most $2j_n$ on which elements of $SL(2, \mathbb{C})$ act by fractional transformations. Using this realization we have (from (1))

$$\begin{aligned} & \bigotimes_n (e^{M_u(\bar{z}_n - z_n)} h_\alpha^{-1}(z_n))_{(n)} \Psi(h_\alpha M_u d\bar{z})(v_n) \\ &= e^{\pi k(u+\alpha)^2/\tau_2} \left(\prod_n e^{-(\alpha - \bar{\alpha})z_n j_n / \tau_2} \right) (2u)^{-J} \\ & \quad \gamma(u+\alpha) \left((e^{(\alpha - \bar{\alpha})z_n / \tau_2} + 2e^{(\alpha - \bar{\alpha})z_n / \tau_2} u v_n) \right). \end{aligned} \quad (5)$$

Now the left hand side is holomorphic in u whereas the right hand side, because of the presence of the u^{-J} factor, is not holomorphic unless

$$\partial_u^{l_0} \partial_{v_1}^{l_1} \dots \partial_{v_N}^{l_N} \gamma(u)(v) \Big|_{\substack{u=\alpha \\ v_i = \exp[(\alpha - \bar{\alpha})z_i / \tau_2]}} = 0 \quad (6)$$

for every $N + 1$ -tuple of non negative integers $L \equiv (l_i)$ such that $|L| \equiv \sum_{i=0}^N l_i < J$.

Conversely, one may see that if (6) holds for $\alpha = 0, 1/2, \tau/2, (\tau + 1)/2$ then Ψ can be extended to the four codimension 1 strata $\mathcal{A}_{(\alpha,0)}^{01}$. Properties of the stratification allow now to apply inductively Hartogs theorem and to extend Ψ , as a holomorphic functional, to all other higher codimension strata. In this way we obtain a quantum state of the theory.

Theorem: There is a one to one correspondence between the space of states of Chern-Simons theory in the presence of Wilson lines and the space $W_N^{fr}(\tau, \underline{z})$ of holomorphic functions $\gamma : \mathbb{C} \rightarrow \bigotimes_n V_{j_n}$ satisfying (4,a-d) and (6) for $\alpha = 0, 1/2, \tau/2, (\tau + 1)/2$.

As an illustration, we will analyze the two simplest cases of zero and one insertions.

States with zero insertions.

From (4,a-d), the zero-points states are simply the even theta-functions of degree $2k$, since for $J = 0$ condition (6) plays no role. The dimension of the space of even theta-functions is $k + 1$. It is spanned by Kac-Moody characters $\chi_{k,j}(\tau, e^{2\pi i u})$ with fixed τ and $j = 0, 1/2, \dots, k/2$.

States with one insertion.

In this case the states with one insertion of spin $j \in \mathbb{N}$ are

$$\gamma(u)(v) = \vartheta(u)v^j,$$

where ϑ is a theta-function of degree $2k$ with

$$\vartheta(-u) = (-1)^j \vartheta(u) \tag{7,a}$$

$$\partial_u^l \vartheta(u)|_{u=\alpha} = 0 \text{ for } l < j \text{ and } \alpha = 0, \frac{1}{2}, \frac{\tau}{2}, \frac{\tau+1}{2}. \tag{7,b}$$

To study the dimension d_j of the space of solutions of (7,a-b) we shall first focus on the case of even spin, $j \in 2\mathbb{N}$. In that case (7,a) requires even theta-functions for which:

$$e^{-4\pi i k u \frac{\alpha - \bar{\alpha}}{\tau - \bar{\tau}}} \vartheta(u + \alpha) = e^{4\pi i k u \frac{\alpha - \bar{\alpha}}{\tau - \bar{\tau}}} \vartheta(-u + \alpha) \quad \alpha = 0, \frac{1}{2}, \frac{\tau}{2}, \frac{\tau+1}{2}. \tag{8}$$

As a consequence of (8), vanishing of $\partial_u^l \vartheta(u)|_{u=\alpha}$ for $l \leq 2n \in 2\mathbb{N}$ implies also vanishing of the same expression for $l = 2n + 1$. Thus we are left with $2j$ linear conditions on the $k + 1$ -dimensional space of even theta-functions and we obtain a lower bound on the dimension of the space of solutions:

$$d_j \geq k - 2j + 1. \tag{9}$$

To obtain an upper bound for d_j we shall use the fact that the sum of multiplicities of zeros of the theta function in any fundamental cell is $2k$. Now, if we have d_j independent even theta-functions with common zeros of total multiplicity $4j \leq 2k$, by a linear combination of them we can obtain another theta-function whose zeros have multiplicity at least $4j + 2(d_j - 1)$. Since the latter has to be $\leq 2k$, it follows that $d_j \geq k - 2j + 1$ and we obtain the expected result

$$d_j = \begin{cases} k - 2j + 1 & \text{for } j \leq k/2, \\ 0 & \text{otherwise.} \end{cases} \tag{10}$$

The case of odd spin j can be treated in a similar way. Now we have $k - 1$ independent odd theta-functions and (7,b) gives $2j - 2$ independent conditions. Finally we obtain the same expression (10) for the dimension.

It can be shown (see [6]) that spaces $W_N^{fr}(\tau, \underline{z})$ form a bundle W_N^{fr} over $\mathbb{H} \times \mathbb{C}^N$. The modular group $\Gamma_N = SL(2, \mathbb{Z}) \ltimes \mathbb{Z}^{2N}$ acts on the base and its action can be lifted to W_N^{fr} . Taking the quotient W_N^{fr}/Γ_N we obtain a new bundle over the moduli space of punctured torus.

Bundle W_N^{fr} is endowed with a flat connection, generalization of the Knizhnik-Zamolodchikov connection [9] to the toroidal geometry [10]. The connection allows to compare states at different points of the base. It is not preserved by the modular group but gives rise to a projective connection on W_N^{fr}/Γ_N .

By studying the behavior of spaces W_N^{fr} on the boundary of the moduli space of punctured tori, we were able to prove the Verlinde formula for their dimensions modulo the result that there are no toroidal states if one of the spins $j_n > k/2$. The latter fact was proven in [4] for the spherical states and in the toroidal case should follow by studying the spherical limit $\tau \rightarrow \infty$ which requires, however, some additional work, the goal of a future research.

References

- [1] Witten, E.: Quantum field theory and the Jones polynomials. *Commun. Math. Phys.* **121**, 351-399 (1989)
- [2] Gawędzki, K.: Wess-Zumino-Witten conformal field theory. In *Constructive Quantum Field Theory II*, eds. A. Wightman and G. Velo, New York: Plenum 1990, p. 89-120
- [3] Elitzur, S., Moore, G., Schwimmer, A., Seiberg, N.: Remarks on the canonical quantization of the Chern-Simons-Witten theory. *Nucl. Phys.* **B326**, 104-134 (1989)
- [4] Gawędzki, K., Kupiainen, A.: $SU(2)$ Chern-Simons theory at genus zero. *Commun. Math. Phys.* **135**, 531-546 (1991)
- [5] Axelrod, S., Della Pietra, S., Witten, E.: Geometric quantization of Chern-Simons gauge theory. *J. Differential Geometry* **33**, 787-902 (1991)
- [6] Falceto, F., Gawędzki, K.: Chern-Simons states at genus one. In preparation.
- [7] Atiyah, M. F., Bott, R.: The Yang-Mills equation over Riemann surfaces. *Phil. Trans. R. Soc. Lond.* **A308**, 523-165 (1982)
- [8] Gunning, R. C.: Lectures on vector bundles over Riemann surfaces. Princeton: Princeton University Press (1967)
- [9] Knizhnik, V., Zamolodchikov, A. B.: Current algebra and Wess-Zumino model in two dimensions. *Nucl. Phys.* **B247**, 83-103 (1984)
- [10] Bernard, D.: On the Wess-Zumino-Witten models on the torus. *Nucl. Phys.* **B303**, 77-93 (1988)